



Scan online to view this article

identifying antibiotic resistance genes in clinical isolates
of *Klebsiella pneumoniae* producing *IMP-1* and *TEM* β -lactamase
Leila Goodarzi¹, Reza Yari¹, Mohsen Mirzaei^{2*}

1-Department of Biology, Borujerd Branch, Islamic Azad University, Borujerd, Iran.

2- Department of Laboratory Sciences, Borujerd Branch, Islamic Azad University, Borujerd, Iran.

Abstract

Aim and Background: *Klebsiella pneumoniae* is a common cause of β -lactamase-associated infections in hospitals. The present study aimed to determine the frequency of antibiotic resistance genes in *Klebsiella pneumoniae* strains producing *IMP-1* and *TEM* β -lactamase.

Materials and methods: The present research identified 94 samples of *K. pneumoniae*, using antibiogram for the phenotypic confirmation of ESBLs. The antibiotic resistance of the isolates and the prevalence of *TEM* and *IMP-1* genes were determined using PCR method.

Findings: Of 94 samples, 77.6% were ESBL-positive and 22.3% ESBL-negative. A total of 4.1% of the samples carried the *IMP-1* gene and 43.8% the *TEM* gene, while 43.8% of the samples carried both genes.

Conclusion: Given that *TEM* and *IMP-1* genes were commonly present in a large number of the resistant samples, physicians are recommended to use therapeutic measures properly, and to prescribe antibiotics rationally.

Keywords: *Klebsiella pneumoniae*, Extended-Spectrum Betalactamase (ESBL), *TEM*, *IMP-1*, Antibiotic Resistance.

Corresponding author:

Department of Laboratory Sciences, Borujerd Branch, Islamic Azad University, Borujerd, Iran.

Email: mirzaei.iaub@gmail.com